

ROD J. ROHRICH, M.D.
PLASTIC SURGERY

FACE/NECK LIFT
AFTER CARE INSTRUCTIONS

1. After surgery elevate your head and shoulders on a 45° angle wedge rest (no pillows!). Don't use any pillows as you want to avoid bending forward at the neck for the first 2 weeks.
2. Avoid straining of any kind for the first 5 days. When awake in bed please flex your ankles and legs up and down 4-6 times every hour.
3. Apply covered ice bags or Swiss eye masks (keep them cold or in ice) to the exposed areas of your face for the first 72 hours to reduce swelling post-operatively.
4. If you have pain or discomfort, take the pain medication every 4-6 hours. It is best to take the pain medication with crackers, jello, etc. If you do not have pain please do not take the pain medication. Do not drink alcohol while you are taking pain medication.
5. A light diet is best following surgery that day. Begin by taking liquid slowly and progress to soups or jello. You may start with a soft, regular diet the **next** day, high in protein.
6. You may have drains in place under the skin, after surgery (in the neck.) These will be secured and you will/can sleep in them – we will show you, but avoid pulling on them. The bulbs connected to the end of the drains should be kept compressed at all times. Instructions regarding the drains will be given to your caretaker. The drains and head dressing are usually removed within 2 days.
7. You can expect some swelling of the face and eyes after surgery. If you develop acute onset of swelling on one side of your face or neck, which is definitely more pronounced than on the other side, or if you are having pain, which is NOT relieved by the pain medication, please call Dr. Rohrich on his cell at 214-500-4870, the UT Southwestern Medical Center office at 214-645-2353, or Dallas Plastic Surgery Institute at 214-821-9114.
8. Avoid turning your head to either side as this will pull on the stitch lines. To turn, move your head and shoulders as a unit.
9. **DO NOT SMOKE.** This is very important.
10. Facial movements (smiling, talking, chewing, yawning, etc.) should be kept to a minimum for the first week.

FACE/NECK LIFT
AFTER CARE INSTRUCTIONS (continued)

11. After your dressings and drains have been removed you may wash your hair with baby shampoo. Lather your hair gently using the palms of your hands and rinse thoroughly until the soap is out. It may take several washings before all the crust and ointment is out of your hair. Do not use hairspray, conditioner, gels, etc. while the stitches and clips are in place. You may start washing your face gently and use a moisturizer on your face, being careful to keep it away from the stitch lines.
12. Your hair may be dried with a blow dryer on a **cool, not a hot**, setting.
13. Hair coloring should be delayed until 3 weeks after surgery when healing is completed and no crust remains.
14. Normal activity can be gradually resumed after 3 weeks but strenuous activity (heart rate above 100 beats per minute and lifting objects greater than 10 pounds) should be avoided for another week. This includes sexual intercourse and sexual activity. The skin of your face is sensitive to sunlight after surgery. Protect your facial skin from excessive exposure to the sun for 8 weeks. Wear a wide brim hat and sunscreen (SPF 20 or greater with UVA and UVB protection) if you are in the sun for prolonged periods, or even on cloudy days for at least 6 months.

GENERAL INSTRUCTIONS AND SCAR CARE:

- Your face/neck will feel tight but not necessarily look tight (it will look natural) and there will be a feeling of numbness in these areas for several weeks to months after surgery. This is normal and will disappear over time and the feeling will return.
- After all sutures have been removed, begin scar treatment: for the first 6 weeks apply a small amount of Eraclea PURE HYDRATION SERUM over all incisions AM & PM. From 6 weeks to 6 months, use Eraclea INTENSIVE REPAIR and RECONSTRUCTION CREAM over all incisions AM & PM as directed by Dr. Rohrich (see scar care instructions). These products can be purchased at both the UT Southwestern and Park Lane locations.
- Cosmetics can be worn on the face as early as 10 days after surgery. However, eyelid or eyelash cosmetics should not be worn until 7 days after all the eyelid sutures have been removed. There may be a feeling of numbness of the eyelids that will subside.
- Take all medications only as directed.

**FACE/NECK LIFT
AFTER CARE INSTRUCTIONS (continued)**

- If you have nausea, vomiting, rash, shortness of breath, or diarrhea after taking your medication, please call the office at (214) 645-2353 or (214) 645-3119.
- If you develop a fever (oral temperature greater than 101°), redness or increased pain at the site of your surgical incisions, please notify Dr. Rohrich immediately.
- **DO NOT WEAR PULLOVER CLOTHING OR HAIRPIECES FOR 3 WEEKS AFTER YOUR FACELIFT, AS YOU COULD DISRUPT YOUR EARLOBE OR HAIRLINE INCISIONS.**
- It is important to be seen by Dr. Rohrich after your initial post-op checks. He will see you in follow-up appointments at 3 weeks, 6 weeks, 6 months and 1 year post-op. Call to schedule your appointments at the UT Southwestern Medical Center office at 214-645-2353 or Dallas Plastic Surgery Institute at 214-821-9114 between the hours of 8:30 – 5:00.

IF YOU HAVE ANY QUESTIONS, PLEASE CALL DR. ROHRICH

If you are a patient at one of the facilities below, please call between the hours of 8:30-5:00. If you need immediate care, please call Dr. Rohrich anytime on his cell. There is no such thing as a minor question. I would rather you call me than wonder and not do the right thing, so please call my office, or email me anytime.

Dr. Rohrich's cell: 214-500-4870
Dr. Rohrich's email: rod.rohrich@utsouthwestern.edu
Dr. Rohrich's website: www.drrohrich.com

Dallas Plastic Surgery Institute
9101 N. Center Expressway, Suite 600
Dallas, TX 75231
214-821-9114

UT Southwestern Medical Center at Dallas
Outpatient Building
1801 Inwood Road
Dallas, TX 75390-9132
214-645-3119